

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD43 (T-Cell Marker) (rSPN/839), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rSPN/839], Mouse, Monoclonal BOT-BNC042220-500**

Article Name	CD43 (T-Cell Marker) (rSPN/839), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rSPN/839], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC042220-500
Supplier Catalog Number	BNC042220-500
Alternative Catalog Number	BOT-BNC042220-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human CD43 protein
Conjugation	CF405S
Product Description	This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43 . Epitope of MAb Bra7G is clearly different from that of MAb rSPN/839, called b as opposed to a for rSPN/839. 70-9...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[rSPN/839]

Molecular Weight	95, 115, or 135 kDa
Isotype	IgG1
UniProt	<a href="#">P16150</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user