

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Human Kappa Light Chain / IGKC (B-Cell Marker) (KLC2289R), CF405S conjugate, 0.1mg/mL, Clone: [KLC2289R], Rabbit, Monoclonal BOT-BNC042289-100**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Human Kappa Light Chain / IGKC (B-Cell Marker) (KLC2289R), CF405S conjugate, 0.1mg/mL, Clone: [KLC2289R], Rabbit, Monoclonal                                                                                                                                  |
| Biozol Catalog Number      | BOT-BNC042289-100                                                                                                                                                                                                                                             |
| Supplier Catalog Number    | BNC042289-100                                                                                                                                                                                                                                                 |
| Alternative Catalog Number | BOT-BNC042289-100-100UL                                                                                                                                                                                                                                       |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Rabbit                                                                                                                                                                                                                                                        |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | IHC                                                                                                                                                                                                                                                           |
| Species Reactivity         | Human                                                                                                                                                                                                                                                         |
| Immunogen                  | Recombinant full-length human Ig kappa chain (IGKC) protein                                                                                                                                                                                                   |
| Conjugation                | CF405S                                                                                                                                                                                                                                                        |
| Product Description        | This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. It recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects al... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [KLC2289R]                                                                                                                                                                                                                                                    |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight  | ~22.5 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| UniProt           | <a href="#">P01601</a>                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Buffer            | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |