

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD44v9 (Marker of Tumor Metastasis) (CD44v9/2344R), CF405S conjugate, 0.1mg/mL, Clone: [CD44v9/2344R], Rabbit, Monoclonal BOT-BNC042344-500**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | CD44v9 (Marker of Tumor Metastasis) (CD44v9/2344R), CF405S conjugate, 0.1mg/mL, Clone: [CD44v9/2344R], Rabbit, Monoclonal                                                                                                                                     |
| Biozol Catalog Number      | BOT-BNC042344-500                                                                                                                                                                                                                                             |
| Supplier Catalog Number    | BNC042344-500                                                                                                                                                                                                                                                 |
| Alternative Catalog Number | BOT-BNC042344-500-500UL                                                                                                                                                                                                                                       |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Rabbit                                                                                                                                                                                                                                                        |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | IHC                                                                                                                                                                                                                                                           |
| Species Reactivity         | Human                                                                                                                                                                                                                                                         |
| Immunogen                  | Recombinant fragment corresponding to the v9 domain of human CD44                                                                                                                                                                                             |
| Conjugation                | CF405S                                                                                                                                                                                                                                                        |
| Product Description        | This antibody recognizes an epitope encoded by exon v4 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemop... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [CD44v9/2344R]                                                                                                                                                                                                                                                |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight  | 80-95 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| UniProt           | <a href="#">P16070</a>                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Buffer            | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |