

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **EGFRvIII (Epidermal Growth Factor Receptor, Variant III) (GFR/2600R), CF405S conjugate, 0.1mg/mL, Clone: [GFR/2600R], Rabbit, Monoclonal BOT-BNC042600-100**

Article Name	EGFRvIII (Epidermal Growth Factor Receptor, Variant III) (GFR/2600R), CF405S conjugate, 0.1mg/mL, Clone: [GFR/2600R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNC042600-100
Supplier Catalog Number	BNC042600-100
Alternative Catalog Number	BOT-BNC042600-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	KLH conjugate of EGFRvIII-specific, 14-amino acid peptide from the predicted amino acid sequence at the fusion junction (LEEKKGNYVVDHC)
Conjugation	CF405S
Product Description	EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor-a (TGF-a), Heparin-binding EGF (HB-EGF), amphiregulin, betacellulin and epi...
Clonality	Monoclonal
Concentration	0.1 mg/mL

Clone Designation	[GFR/2600R]
Molecular Weight	~145 kDa (vIII variant)
UniProt	<a href="#">P00533</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user