

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Uroplakin 1A (Urothelial Differentiation Marker)(UPK1A/2924), CF405S conjugate, 0.1mg/mL, Clone: [UPK1A/2924], Mouse, Monoclonal BOT-BNC042924-100

Article Name	Uroplakin 1A (Urothelial Differentiation Marker)(UPK1A/2924), CF405S conjugate, 0.1mg/mL, Clone: [UPK1A/2924], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC042924-100
Supplier Catalog Number	BNC042924-100
Alternative Catalog Number	BOT-BNC042924-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa 114-173) of human Uroplakin 1A (UPK1A) protein (exact sequence is proprietary)
Conjugation	CF405S
Product Description	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins me...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[UPK1A/2924]

Molecular Weight	29 kDa
UniProt	O00322
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry