

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin, pan (Epithelial Marker)(PCK/3150), CF405S conjugate, 0.1mg/mL, Clone: [PCK/3150], Mouse, Monoclonal BOT-BNC043150-500

Article Name	Cytokeratin, pan (Epithelial Marker)(PCK/3150), CF405S conjugate, 0.1mg/mL, Clone: [PCK/3150], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC043150-500
Supplier Catalog Number	BNC043150-500
Alternative Catalog Number	BOT-BNC043150-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Canine, Feline, Human
Immunogen	Crude cytokeratin extract prepared from RT-4 and MCF-7 cells.
Conjugation	CF405S
Product Description	Anti-cytokeratin clone PCK/3150 demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, PCK/3150 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial ep...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[PCK/3150]

Molecular Weight	40-67 kDa (Multiple)
UniProt	Multiple
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry