

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 8 (KRT8) (KRT8/2174R), CF640R conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal BOT-BNC402174-500

| | |
|----------------------------|---|
| Article Name | Cytokeratin 8 (KRT8) (KRT8/2174R), CF640R conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC402174-500 |
| Supplier Catalog Number | BNC402174-500 |
| Alternative Catalog Number | BOT-BNC402174-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Bovine, Human, Porcine, Rabbit, Sheep |
| Immunogen | Recombinant full-length human KRT8 protein |
| Conjugation | CF640R |
| Product Description | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [KRT8/2174R] |

| | |
|-------------------|--|
| Molecular Weight | 52.5 kDa |
| UniProt | P05787 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |