

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Thyroglobulin(2H11 + 6E1), CF647 conjugate, 0.1mg/mL, Clone: [2H11 6E1], Mouse, Monoclonal BOT-BNC470724-500

| | |
|----------------------------|--|
| Article Name | Thyroglobulin(2H11 + 6E1), CF647 conjugate, 0.1mg/mL, Clone: [2H11 6E1], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC470724-500 |
| Supplier Catalog Number | BNC470724-500 |
| Alternative Catalog Number | BOT-BNC470724-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Human thyroid follicular cells |
| Conjugation | CF647 |
| Product Description | Thyroglobulin is a 660 kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo c... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [2H11 6E1] |

| | |
|-------------------|---|
| Molecular Weight | 660 kDa (Dimeric Form) |
| UniProt | P01266 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1.0 ug/million cells in 0.1 mL Optimal dilution for a specific application should be determined by user |