

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD20 / MS4A1 (B-Cell Marker)(L26 + IGEL/773), CF647 conjugate, 0.1mg/mL, Clone: [L26 IGEL/773], Mouse, Monoclonal BOT-BNC470854-500**

Article Name	CD20 / MS4A1 (B-Cell Marker)(L26 + IGEL/773), CF647 conjugate, 0.1mg/mL, Clone: [L26 IGEL/773], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC470854-500
Supplier Catalog Number	BNC470854-500
Alternative Catalog Number	BOT-BNC470854-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human, Monkey, Primate
Immunogen	Human tonsil B cells (L26), Recombinant human MS4A1 protein (IGEL/773)
Conjugation	CF647
Product Description	This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[L26 IGEL/773]
Molecular Weight	33-37 kDa

UniProt	<a href="#">P11836</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Does not react with Cow, Dog, Pig or Rat For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry