

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

WT1 (Wilms Tumor 1)(WT1/857), CF647 conjugate, 0.1mg/mL, Clone: [WT1/857], Mouse, Monoclonal BOT-BNC470857-100

Article Name	WT1 (Wilms Tumor 1)(WT1/857), CF647 conjugate, 0.1mg/mL, Clone: [WT1/857], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC470857-100
Supplier Catalog Number	BNC470857-100
Alternative Catalog Number	BOT-BNC470857-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Mouse, Rat
Immunogen	Recombinant human WT1 protein
Conjugation	CF647
Product Description	Recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[WT1/857]

Molecular Weight	47-55 kDa
UniProt	P19544
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user