

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Myeloid-Associated Differentiation Marker(MYADM/971), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [MYADM/971], Mouse, Monoclonal BOT-BNC470971-500

| | |
|----------------------------|---|
| Article Name | Myeloid-Associated Differentiation Marker(MYADM/971), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [MYADM/971], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC470971-500 |
| Supplier Catalog Number | BNC470971-500 |
| Alternative Catalog Number | BOT-BNC470971-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Primate |
| Immunogen | Recombinant human MYADM protein |
| Conjugation | CF647 |
| Product Description | Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellul... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [MYADM/971] |

| | |
|-------------------|--|
| Molecular Weight | Unknown |
| Isotype | IgG1 |
| UniProt | Q96S97 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |