

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD3e(CRIS-7), CF647 conjugate, 0.1mg/mL, Clone: [CRIS-7], Mouse, Monoclonal BOT-BNC471050-500**

|                            |   |
|----------------------------|---|
| Article Name               | CD3e(CRIS-7), CF647 conjugate, 0.1mg/mL, Clone: [CRIS-7], Mouse, Monoclonal   |
| Biozol Catalog Number      | BOT-BNC471050-500   |
| Supplier Catalog Number    | BNC471050-500   |
| Alternative Catalog Number | BOT-BNC471050-500-500UL   |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC, Functional Studies, IF  |
| Species Reactivity         | Human, Monkey   |
| Immunogen                  | Stimulated human leukocytes   |
| Conjugation                | CF647   |
| Product Description        | Recognizes the epsilon chain of CD3 (Workshop V, Code: CD03.09), which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closely associated at the lym... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [CRIS-7]  |

|                   |  |
|-------------------|--|
| Molecular Weight  | 20 kDa   |
| UniProt           | <a href="#">P07766</a>   |
| Buffer            | PBS, 0.1% BSA, 0.05% azide   |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Induce T cell activation and proliferation Optimal dilution for a specific application should be determined by user |