

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human Papillomavirus 16 (HPV-16)(HPV16/1058), CF647 conjugate, 0.1mg/mL, Clone: [HPV16/1058], Mouse, Monoclonal BOT-BNC471058-500

| | |
|----------------------------|---|
| Article Name | Human Papillomavirus 16 (HPV-16)(HPV16/1058), CF647 conjugate, 0.1mg/mL, Clone: [HPV16/1058], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC471058-500 |
| Supplier Catalog Number | BNC471058-500 |
| Alternative Catalog Number | BOT-BNC471058-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Virus |
| Immunogen | Human papilloma virus type 16, major capsid protein L1 |
| Conjugation | CF647 |
| Product Description | Reacts with a protein of 57 kDa, identified as the L1 protein of human papilloma virus type 16 (HPV-16). It is the major capsid protein of HPV-16. Infection with specific types of HPV has been associated with an increased risk of developing cervical ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [HPV16/1058] |

| | |
|-------------------|---|
| Molecular Weight | 57 kDa |
| UniProt | Not Applicable |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-paraffin 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1.0 ug/mL Immunoprecipitation 0.5-1 ug/500 ug protein lysate Optimal dilution for a specific application should be determined by user |