

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Catenin, beta (p120) (15B8), CF647 conjugate, 0.1mg/mL, Clone: [15B8], Mouse, Monoclonal BOT-BNC471443-500**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Catenin, beta (p120) (15B8), CF647 conjugate, 0.1mg/mL, Clone: [15B8], Mouse, Monoclonal                                                                                                                                                                      |
| Biozol Catalog Number      | BOT-BNC471443-500                                                                                                                                                                                                                                             |
| Supplier Catalog Number    | BNC471443-500                                                                                                                                                                                                                                                 |
| Alternative Catalog Number | BOT-BNC471443-500-500UL                                                                                                                                                                                                                                       |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Mouse                                                                                                                                                                                                                                                         |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | IF, IHC, WB                                                                                                                                                                                                                                                   |
| Species Reactivity         | Bovine, Canine, Gallus, Human, Mouse, Rat                                                                                                                                                                                                                     |
| Immunogen                  | Recombinant chicken $\gamma$ -catenin                                                                                                                                                                                                                         |
| Conjugation                | CF647                                                                                                                                                                                                                                                         |
| Product Description        | Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [15B8]                                                                                                                                                                                                                                                        |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|-------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight  | 92 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| UniProt           | <a href="#">P35222</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Buffer            | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |