

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Human IgM Immunoglobulin(ICO-30), CF568 conjugate, 0.1mg/mL, Clone: [ICO-30], Mouse, Monoclonal BOT-BNC680364-100**

Article Name	Human IgM Immunoglobulin(ICO-30), CF568 conjugate, 0.1mg/mL, Clone: [ICO-30], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC680364-100
Supplier Catalog Number	BNC680364-100
Alternative Catalog Number	BOT-BNC680364-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Heavy chain of human IgM
Conjugation	CF568
Product Description	Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. Monomeric Ig...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[ICO-30]

Molecular Weight	50-75 kDa
UniProt	<a href="#">P01871</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user