

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

p40(Rabbit PAb), CF568 conjugate, 0.1mg/mL, Clone: [PAb], Monoclonal BOT-BNC680375-100

Article Name	p40(Rabbit PAb), CF568 conjugate, 0.1mg/mL, Clone: [PAb], Monoclonal
Biozol Catalog Number	BOT-BNC680375-100
Supplier Catalog Number	BNC680375-100
Alternative Catalog Number	BOT-BNC680375-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC, WB
Species Reactivity	Bovine, Human, Mouse, Rat
Immunogen	A synthetic peptide (ENNAQTQFSEPQY) corresponding to aa5-17 of human p40
Conjugation	CF568
Product Description	p40 (p63 delta) is a marker determined to be highly specific for squamous basal cells by immunohistochemistry (IHC). The current more routinely recommended marker, p63, appears to have less specificity compared to p40, especially on squamous cell tum...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[PAb]

Molecular Weight	40 kDa
UniProt	Q9H3D4
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user