

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Napsin-A(NAPSA/1239), CF568 conjugate, 0.1mg/mL, Clone: [NAPSA/1239], Mouse, Monoclonal BOT-BNC681239-100

| | |
|----------------------------|--|
| Article Name | Napsin-A(NAPSA/1239), CF568 conjugate, 0.1mg/mL, Clone: [NAPSA/1239], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC681239-100 |
| Supplier Catalog Number | BNC681239-100 |
| Alternative Catalog Number | BOT-BNC681239-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human Napsin-A protein |
| Conjugation | CF568 |
| Product Description | Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high exp... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [NAPSA/1239] |

| | |
|-------------------|--|
| Molecular Weight | 37 kDa |
| UniProt | O96009 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |