

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) (KRTL/1577R), CF568 conjugate, 0.1mg/mL, Clone: [KRTL/1577R], Rabbit, Monoclonal BOT-BNC681577-100

Article Name	Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) (KRTL/1577R), CF568 conjugate, 0.1mg/mL, Clone: [KRTL/1577R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNC681577-100
Supplier Catalog Number	BNC681577-100
Alternative Catalog Number	BOT-BNC681577-100-100UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human KRT77 protein fragment
Conjugation	CF568
Product Description	This MAb recognizes the 56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI <5.7) an...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[KRTL/1577R]

Molecular Weight	56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19)
UniProt	Q7Z794
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user