

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD62L (L-Selectin)(CD62L/1588), CF568 conjugate, 0.1mg/mL, Clone: [CD62L/1588], Mouse, Monoclonal BOT-BNC681588-500

| | |
|----------------------------|---|
| Article Name | CD62L (L-Selectin)(CD62L/1588), CF568 conjugate, 0.1mg/mL, Clone: [CD62L/1588], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC681588-500 |
| Supplier Catalog Number | BNC681588-500 |
| Alternative Catalog Number | BOT-BNC681588-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | MAB raised against supernatant from phorbol myristic acid activated human peripheral blood leukocytes. |
| Conjugation | CF568 |
| Product Description | Selectins, also designated CD62 antigens, comprise a family of carbohydrate-binding proteins involved in mediating cellular interactions with leukocytes. L-Selectin (also designated LECAM-1 or CD62L) is expressed on the majority of B and naive T cell... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [CD62L/1588] |
| Molecular Weight | 65-75 kDa |

| | |
|-------------------|--|
| UniProt | P14151 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |