

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Factor XIIIa (Coagulation Factor XIIIa Chain) (F13A1/1683), CF568 conjugate, 0.1mg/mL, Clone: [F13A1/1683], Mouse, Monoclonal BOT-BNC681683-500

| | |
|----------------------------|---|
| Article Name | Factor XIIIa (Coagulation Factor XIIIa Chain) (F13A1/1683), CF568 conjugate, 0.1mg/mL, Clone: [F13A1/1683], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC681683-500 |
| Supplier Catalog Number | BNC681683-500 |
| Alternative Catalog Number | BOT-BNC681683-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is proprietary) |
| Conjugation | CF568 |
| Product Description | This antibody recognizes a protein of 83 kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macropha... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [F13A1/1683] |

| | |
|-------------------|--|
| Molecular Weight | 83 kDa |
| UniProt | P00488 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user |