

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (F8/86), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [F8/86], Mouse, Monoclonal BOT-BNC681716-500**

Article Name	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (F8/86), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [F8/86], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC681716-500
Supplier Catalog Number	BNC681716-500
Alternative Catalog Number	BOT-BNC681716-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Factor isolated from human plasma von Willebr
Conjugation	CF568
Product Description	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[F8/86]

Molecular Weight	250 kDa
Isotype	IgG1
UniProt	<a href="#">P04275</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>