

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Human Immunoglobulin Gamma (IgG) Heavy Chain(IG217 + IG266), CF488A conjugate, 0.1mg/mL, Clone: [IG217 IG266], Mouse, Monoclonal BOT-BNC881262-500

Article Name	Human Immunoglobulin Gamma (IgG) Heavy Chain(IG217 + IG266), CF488A conjugate, 0.1mg/mL, Clone: [IG217 IG266], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC881262-500
Supplier Catalog Number	BNC881262-500
Alternative Catalog Number	BOT-BNC881262-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Purified human IgG
Conjugation	CF488A
Product Description	This MAb is specific to heavy chain of IgG and shows minimal cross-reaction with heavy chains of other immunoglobulins. It is reactive with all subclasses of Gamma heavy chain(IgG1, IgG2a, IgG2b, IgG3). The most common feature of plasmacytomas, and c...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[IG217 IG266]

Molecular Weight	75 kDa
UniProt	P01857
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Flow cytometry: 1 ug/million cells Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user