

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 7 (Glandular and Transitional Epithelial Marker) (KRT7/1499R), CF488A conjugate, 0.1mg/mL, Clone: [KRT7/1499R], Rabbit, Monoclonal BOT-BNC881499-500

| | |
|----------------------------|---|
| Article Name | Cytokeratin 7 (Glandular and Transitional Epithelial Marker) (KRT7/1499R), CF488A conjugate, 0.1mg/mL, Clone: [KRT7/1499R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC881499-500 |
| Supplier Catalog Number | BNC881499-500 |
| Alternative Catalog Number | BOT-BNC881499-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human KRT7 protein |
| Conjugation | CF488A |
| Product Description | This antibody recognizes an intermediate filament protein (IFP) of 55 kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [KRT7/1499R] |

| | |
|-------------------|--|
| Molecular Weight | 55 kDa |
| UniProt | P08729 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |