

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Blood Group Antigen A(HE-193), 0.2mg/mL, Clone: [HE-193], Mouse, Monoclonal BOT-BNUB1064-500

| | |
|----------------------------|---|
| Article Name | Blood Group Antigen A(HE-193), 0.2mg/mL, Clone: [HE-193], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1064-500 |
| Supplier Catalog Number | BNUB1064-500 |
| Alternative Catalog Number | BOT-BNUB1064-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Mixture of erythrocytes of blood group A and glycoprotein fraction isolated from the saliva of secretors with blood group A |
| Product Description | This antibody is applicable for staining ABO blood group A (monofucosyl and difucosyl A antigens with chain types 1, 2, 3, 4, 5, 6). It also recognizes Forssman antigen. The histo-blood group ABO involves three carbohydrate antigens: A, B, and H. Blo... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [HE-193] |
| Molecular Weight | Multiple |

| | |
|-------------------|---|
| UniProt | P16442 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin/paraffin tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |