

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CDw17(H018.3G-6.F5), 0.2mg/mL, IgM, Clone: [H018.3G-6.F5], Mouse, Monoclonal BOT-BNUB1092-100

Article Name	CDw17(H018.3G-6.F5), 0.2mg/mL, IgM, Clone: [H018.3G-6.F5], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1092-100
Supplier Catalog Number	BNUB1092-100
Alternative Catalog Number	BOT-BNUB1092-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	?-2 Microglobulin associated proteins from a detergent lysate of human PBL
Product Description	CDw17 / CD17 is an intermediate glycosphingolipid from the metabolism of higher gangliosides that localizes to sphingolipid-sterol rafts. CD17 is detectable in monocytes, granulocytes, basophils, platelets, a subset of peripheral B cells (CD19) and ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[H018.3G-6.F5]
Molecular Weight	Not Known
Isotype	IgM

UniProt	Not Known
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry