

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) (KRTH/1576R), 0.2mg/mL, Clone: [KRTH/1576R], Rabbit, Monoclonal BOT-BNUB1576-500

Article Name	Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) (KRTH/1576R), 0.2mg/mL, Clone: [KRTH/1576R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB1576-500
Supplier Catalog Number	BNUB1576-500
Alternative Catalog Number	BOT-BNUB1576-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human KRT76 protein
Product Description	This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67 kDa (CK1), 64 kDa (CK3), 59 kDa (CK4), 58 kDa (CK5), 56 kDa (CK6), 52 kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[KRTH/1576R]
Molecular Weight	52-67 kDa

UniProt	Q01546
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user