

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD79a (IGA/1688R), 0.2mg/mL, Clone: [IGA/1688R], Rabbit, Monoclonal BOT-BNUB1688-500

Article Name	CD79a (IGA/1688R), 0.2mg/mL, Clone: [IGA/1688R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB1688-500
Supplier Catalog Number	BNUB1688-500
Alternative Catalog Number	BOT-BNUB1688-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein.
Product Description	A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[IGA/1688R]
Molecular Weight	44 kDa

UniProt	P11912
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user