

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

p53 Tumor Suppressor Protein (rBP53-12), 0.2mg/mL, IgG1, Clone: [rBP53-12], Mouse, Monoclonal BOT-BNUB2094-100

Article Name	p53 Tumor Suppressor Protein (rBP53-12), 0.2mg/mL, IgG1, Clone: [rBP53-12], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB2094-100
Supplier Catalog Number	BNUB2094-100
Alternative Catalog Number	BOT-BNUB2094-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Canine, Gallus, Hamster, Human, Monkey
Immunogen	Recombinant human wild-type p53 protein
Product Description	This MAb reacts with an N-terminal epitope (aa 16-25) of both wild type and mutated p53. Mutation and/or allelic loss of p53 is one of the causes of a variety of mesenchymal and epithelial tumors. If it occurs in the germ line, such tumors run in fam...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[rBP53-12]
Molecular Weight	53 kDa

Isotype	IgG1
UniProt	P04637
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user