

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 19(A53-B/A2.26), 1mg/mL, Clone: [A53-B/A2.26], Mouse, Monoclonal BOT-BNUM0020-50

Article Name	Cytokeratin 19(A53-B/A2.26), 1mg/mL, Clone: [A53-B/A2.26], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM0020-50
Supplier Catalog Number	BNUM0020-50
Alternative Catalog Number	BOT-BNUM0020-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IF, IHC, WB
Species Reactivity	Human
Immunogen	Human breast cancer MCF-7 cells
Product Description	This MAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40 kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, ...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[A53-B/A2.26]
Molecular Weight	40 kDa

UniProt	P08727
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat or ferret, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user