

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD28(C28/74), 1mg/mL, Clone: [C28/74], Mouse, Monoclonal BOT-BNUM0074-50

| | |
|----------------------------|---|
| Article Name | CD28(C28/74), 1mg/mL, Clone: [C28/74], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM0074-50 |
| Supplier Catalog Number | BNUM0074-50 |
| Alternative Catalog Number | BOT-BNUM0074-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Recombinant human CD28 protein |
| Product Description | Recognizes a glycoprotein of 44-88 kDa, which is identified as CD28. It is the critical T-cell co-stimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [C28/74] |
| Molecular Weight | 44 kDa (monomer), 88 kDa (dimer) |
| UniProt | P10747 |
| Buffer | PBS, no BSA, no azide |

| | |
|-------------------|--|
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |