

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TTF-1 / NKX2.1(NX2.1/690), 1mg/mL, Clone: [NX2.1/690], Mouse, Monoclonal BOT-BNUM0690-50

| | |
|----------------------------|---|
| Article Name | TTF-1 / NKX2.1(NX2.1/690), 1mg/mL, Clone: [NX2.1/690], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM0690-50 |
| Supplier Catalog Number | BNUM0690-50 |
| Alternative Catalog Number | BOT-BNUM0690-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Recombinant TTF-1 protein |
| Product Description | Recognizes a protein of 40 kDa, identified as Thyroid transcription factor-1 (TTF-1). TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver,... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [NX2.1/690] |
| Molecular Weight | 40 kDa |

| | |
|-------------------|--|
| UniProt | P43699 |
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |