

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mucin 1 / EMA / Episialin / CD227(GP1.4 + E29), 1mg/mL, Clone: [GP1.4 E29], Mouse, Monoclonal BOT-BNUM0743-50

| | |
|----------------------------|---|
| Article Name | Mucin 1 / EMA / Episialin / CD227(GP1.4 + E29), 1mg/mL, Clone: [GP1.4 E29], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM0743-50 |
| Supplier Catalog Number | BNUM0743-50 |
| Alternative Catalog Number | BOT-BNUM0743-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Human milk fat globule membranes (GP1.4), Delipidated extract of human milk fat globule membranes (E29) |
| Product Description | In Western blotting, it recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protecti... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [GP1.4 E29] |
| Molecular Weight | 265-400 kDa |

| | |
|-------------------|--|
| UniProt | P15941 |
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |