

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **HLA-DRB(HLA-DRB/1067), 1mg/mL, Clone: [HLA-DRB/1067], Mouse, Monoclonal BOT-BNUM1067-50**

Article Name	HLA-DRB(HLA-DRB/1067), 1mg/mL, Clone: [HLA-DRB/1067], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM1067-50
Supplier Catalog Number	BNUM1067-50
Alternative Catalog Number	BOT-BNUM1067-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human, Monkey
Immunogen	Activated human peripheral blood mononuclear cells
Product Description	This MAb reacts with the beta-chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. Its epitope is different from that of MAb L243. HLA-DR is a heterodimeric cell surface glycoprotein comprised...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[HLA-DRB/1067]
Molecular Weight	28 kDa (beta chain)

UniProt	<a href="#">P01911</a>
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user