

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) (Ki-1/1747R), 1mg/mL, Clone: [Ki-1/1747R], Rabbit, Monoclonal BOT-BNUM1747-50

Article Name	CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) (Ki-1/1747R), 1mg/mL, Clone: [Ki-1/1747R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUM1747-50
Supplier Catalog Number	BNUM1747-50
Alternative Catalog Number	BOT-BNUM1747-50-50UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human full-length TNFRSF8 protein
Product Description	This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In ...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[Ki-1/1747R]
Molecular Weight	105-120 kDa

UniProt	P28908
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 8.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user