

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bcl-X (Apoptosis Marker)(BCL2L1/1762), 1mg/mL, Clone: [BCL2L1/1762], Mouse, Monoclonal BOT-BNUM1762-50

Article Name	Bcl-X (Apoptosis Marker)(BCL2L1/1762), 1mg/mL, Clone: [BCL2L1/1762], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM1762-50
Supplier Catalog Number	BNUM1762-50
Alternative Catalog Number	BOT-BNUM1762-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant human BCL2L1 protein fragment (around aa 1-135) (exact sequence is proprietary)
Product Description	Bcl-x belongs to the Bcl-2 family of proteins and plays a crucial role in apoptosis. Bcl-X has two isoforms: Bcl-XL (long), a 26 kDa protein which suppresses cell death, and Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain whic...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[BCL2L1/1762]
Molecular Weight	27 kDa
UniProt	Q07817

Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry