

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), 1mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal BOT-BNUM2034-50

Article Name	CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), 1mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUM2034-50
Supplier Catalog Number	BNUM2034-50
Alternative Catalog Number	BOT-BNUM2034-50-50UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human CD22 protein (exact sequence is proprietary)
Product Description	This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[LPFS2/2034R]
Molecular Weight	120 kDa

UniProt	P26842
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user