

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cyclin D2(CCND2/2620), 1mg/mL, Clone: [CCND2/2620], Mouse, Monoclonal BOT-BNUM2620-50

| | |
|----------------------------|---|
| Article Name | Cyclin D2(CCND2/2620), 1mg/mL, Clone: [CCND2/2620], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM2620-50 |
| Supplier Catalog Number | BNUM2620-50 |
| Alternative Catalog Number | BOT-BNUM2620-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human Cyclin D2 (CCND2) protein |
| Product Description | Cyclins are a family of proteins that control how cells proceed through the multi-step cycle of cell division. Cyclin D2 helps to regulate a step in the cycle called the G1-S transition, in which the cell moves from the G1 phase, when cell growth occ... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [CCND2/2620] |
| Molecular Weight | 34 kDa |
| UniProt | P30279 |

| | |
|-------------------|--|
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |