

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), 1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal BOT-BNUM2651-50

Article Name	Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), 1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUM2651-50
Supplier Catalog Number	BNUM2651-50
Alternative Catalog Number	BOT-BNUM2651-50-50UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human Interleukin 10 (IL10) protein
Product Description	Interleukin 10 (IL10) is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immuno-regulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II an...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[IL10/2651R]
Molecular Weight	20 kDa (Monomer), 37 kDa (Dimer)

UniProt	P22301
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user