

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Langerin / CD207 (Marker of Langerhans Cells)(LGRN/3136R), CF740 conjugate, 0.1mg/mL, Clone: [LGRN/3136R], CF 740, Rabbit, Monoclonal BOT-BNC743136-500**

Article Name	Langerin / CD207 (Marker of Langerhans Cells)(LGRN/3136R), CF740 conjugate, 0.1mg/mL, Clone: [LGRN/3136R], CF 740, Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNC743136-500
Supplier Catalog Number	BNC743136-500
Alternative Catalog Number	BOT-BNC743136-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa74-213) of human Langerin protein (exact sequence is proprietary)
Conjugation	CF 740
Product Description	Langerhans cells (LCs) are a subset of immature dendritic cells (DCs) that specifically localize in the epidermis and other mucosal epithelia. Epidermal LCs possess strong immuno-stimulatory capacity and play a central role in the initiation and regu...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[LGRN/3136R]

Molecular Weight	48 kDa
UniProt	<a href="#">Q9UJ71</a>
Buffer	PBS, 0.1% rBSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry