

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Bcl-6 (Follicular Lymphoma Marker)(rBCL6/1475), CF740 conjugate, 0.1mg/mL, Clone: [rBCL6/1475], CF 740, Mouse, Monoclonal BOT-BNC743433-100**

Article Name	Bcl-6 (Follicular Lymphoma Marker)(rBCL6/1475), CF740 conjugate, 0.1mg/mL, Clone: [rBCL6/1475], CF 740, Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC743433-100
Supplier Catalog Number	BNC743433-100
Alternative Catalog Number	BOT-BNC743433-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant human bcl-6 protein fragment (around aa256-389) (Exact sequence is proprietary)
Conjugation	CF 740
Product Description	Bcl-6 is a transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different core...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[rBCL6/1475]
Molecular Weight	95 kDa

UniProt	<a href="#">P41182</a>
Buffer	PBS, 0.1% rBSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry