

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Biotinylated Human Mesothelin (C-6His-Avi) EBT-EPT283

Article Name	Biotinylated Human Mesothelin (C-6His-Avi)
Biozol Catalog Number	EBT-EPT283
Supplier Catalog Number	EPT283
Alternative Catalog Number	EBT-EPT283-100
Manufacturer	ELK Biotechnology
Category	Proteine/Peptide
Product Description	Biotinylated Recombinant Human Mesothelin is produced by our Mammalian expression system and the target gene encoding Glu296-Ser598 is expressed with a 6His, Avi tag at the C-terminus....
Molecular Weight	Molecular weight: 36.7 KDa. Apparent molecular weight: 38-60 KDa, reducing conditions
UniProt	AAH09272.1
Purity	Greater than 95% as determined by reducing SDS-PAGE.

Application Notes	Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test. Biological activity: Loaded Anti-Human Mesothelin mAb-Fc(CatNC078) on Protein A Biosensor, can bind Biotinylated Human Mesothelin-His-Avi(CatCY05) with an affinity constant of 10⁻³ nM as determined in BLI assay. Background: Mesothelin is a cell surface glycoprotein whose expression is limited to mesothelial cells of the serosa (pleura, pericardium, and peritoneum) and epithelial cells of the trachea, tonsils, fallopian tube, and kidneys. Mesothelin plays an important role in cell survival, proliferation, migration, invasion, tumor progression, and resistance to chemotherapy. The overexpression of mesothelin can activate NF-kappaB and signal transducer and activator of transcription 3 (Stat3), inhibit apoptotic signaling and TNF-alpha-induced apoptosis, and accelerate the G1-S transition. Mesothelin is also found overexpressed in various cancers, including malignant mesothelioma, pancreatic or ovarian carcinoma, sarcomas and in some gastrointestinal or pulmonary carcinomas. As a result of its limited expression in normal tissues, mesothelin has been reported as an ideal tumor-associated marker for the development of targeted therapy
-------------------	---