

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Monoclonal Antibody to HLA-DRA (MHC II)(Clone : 19-26.1, same as MB-26.1), Clone: [19-26.1, same as MB-26.1], Mouse ABI-36-1231**

|                            |   |
|----------------------------|---|
| Article Name               | Monoclonal Antibody to HLA-DRA (MHC II)(Clone : 19-26.1, same as MB-26.1), Clone: [19-26.1, same as MB-26.1], Mouse   |
| Biozol Catalog Number      | ABI-36-1231   |
| Supplier Catalog Number    | 36-1231   |
| Alternative Catalog Number | ABI-36-1231-100UG   |
| Manufacturer               | Abeomics  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FACS, IF, IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Raji Cells  |
| Product Description        | This MAb reacts with the HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. ... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [19-26.1, same as MB-26.1]  |
| NCBI                       | <a href="#">3122</a>  |
| UniProt                    | <a href="#">P01903</a>  |

|                    |   |
|--------------------|---|
| Purity             | Affinity Chromatography   |
| Form               | Purified  |
| Target             | HLA-DRA (MHC II)  |
| Application Dilute | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min a |