

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Monoclonal Antibody to Cytokeratin 17 (KRT17) (Basal Epithelial Marker)(Clone : E3, same as Ks17.E3), Clone: [E3, same as Ks17.E3], Mouse ABI-36-1370

| | |
|----------------------------|---|
| Article Name | Monoclonal Antibody to Cytokeratin 17 (KRT17) (Basal Epithelial Marker)(Clone : E3, same as Ks17.E3), Clone: [E3, same as Ks17.E3], Mouse |
| Biozol Catalog Number | ABI-36-1370 |
| Supplier Catalog Number | 36-1370 |
| Alternative Catalog Number | ABI-36-1370-100UG |
| Manufacturer | Abeomics |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC, WB |
| Species Reactivity | Human, Rat |
| Immunogen | The cytoskeletal fraction of Rat colon epithelium |
| Product Description | Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ... |
| Clonality | Monoclonal |
| Clone Designation | [E3, same as Ks17.E3] |
| NCBI | 3872 |

| | |
|--------------------|---|
| UniProt | Q04695 |
| Purity | Affinity Chromatography |
| Form | Purified |
| Target | Cytokeratin 17 (KRT17) (Basal Epithelial Marker) |
| Application Dilute | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Frozen & formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris w |