

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Monoclonal Antibody to Bcl-2 (Apoptosis & Follicular Lymphoma Marker)(Clone : 8C8), Clone: [8C8], Mouse ABI-36-1641

| | |
|----------------------------|---|
| Article Name | Monoclonal Antibody to Bcl-2 (Apoptosis & Follicular Lymphoma Marker)(Clone : 8C8), Clone: [8C8], Mouse |
| Biozol Catalog Number | ABI-36-1641 |
| Supplier Catalog Number | 36-1641 |
| Alternative Catalog Number | ABI-36-1641-100UG |
| Manufacturer | Abeomics |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein. |
| Product Description | This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 i oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 i oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphoma... |
| Clonality | Monoclonal |
| Clone Designation | [8C8] |
| NCBI | 596 |
| UniProt | P10415 |

| | |
|--------------------|---|
| Purity | Affinity Chromatography |
| Form | Purified |
| Target | Bcl-2 (Apoptosis & Follicular Lymphoma Marker) |
| Application Dilute | Flow Cytometry (1-2ug/million cells), ,Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1m |