

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Monoclonal Antibody to von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(Clone : SPM577), Clone: [SPM577], Mouse ABI-36-1805

| | |
|----------------------------|---|
| Article Name | Monoclonal Antibody to von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(Clone : SPM577), Clone: [SPM577], Mouse |
| Biozol Catalog Number | ABI-36-1805 |
| Supplier Catalog Number | 36-1805 |
| Alternative Catalog Number | ABI-36-1805-100UG |
| Manufacturer | Abeomics |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human vWF fragment |
| Product Description | von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation... |
| Clonality | Monoclonal |
| Clone Designation | [SPM577] |
| NCBI | 7450 |
| UniProt | P04275 |

| | |
|--------------------|--|
| Purity | Affinity Chromatography |
| Form | Purified |
| Target | von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) |
| Application Dilute | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM |