

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Fab anti-Human IgG (H+L)-FITC, MinX none DNA-SEC-183948

Article Name	Goat Fab anti-Human IgG (H+L)-FITC, MinX none
Biozol Catalog Number	DNA-SEC-183948
Supplier Catalog Number	SEC-183948
Alternative Catalog Number	DNA-SEC-183948
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	FITC
Format	Fab
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Fab Anti-Human IgG (H&L) Fluorescein Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum imm...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.07, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Fab Anti-Human IgG (H&L) Fluorescein Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.